

On Zumkeller Numbers

K.P.S. Bhaskara Rao Yuejian Peng

Department of Mathematics and Computer Science

Indiana State University

Terre Haute, IN, 47809, USA

Email: bkopparty@indstate.edu; yuejian.peng@indstate.edu

Abstract

Generalizing the concept of a perfect number, Sloane's sequences of integers A083207 lists the sequence of integers n with the property: the positive factors of n can be partitioned into two disjoint parts so that the sums of the two parts are equal. Following [4] Clark et al., we shall call such integers, Zumkeller numbers. Generalizing this, in [4] a number n is called a half-Zumkeller number if the positive proper factors of n can be partitioned into two disjoint parts so that the sums of the two parts are equal.

An extensive study of properties of Zumkeller numbers, half-Zumkeller numbers and their relation to practical numbers is undertaken in this paper.

In [4] Clark et al., announced results about Zumkellers numbers and half-Zumkeller numbers and suggested two conjectures. In the present paper we shall settle one of the conjectures, prove the second conjecture in some special cases and prove several results related to the second conjecture. We shall also show that if there is an even Zumkeller number that is not half-Zumkeller it should be bigger than 7.2334989×10^9 .

1 Introduction

A positive integer n is called a *perfect* number if n equals the sum of its proper positive factors.

Generalizing this concept in 2003, Zumkeller published in Sloane's sequences of integers A083207 a sequence of integers n with the property that the positive factors of n

can be partitioned into two disjoint parts so that the sums of the two parts are equal.

Following Clark et al., we shall call such integers, Zumkeller numbers. In [1] Clark et al., announced several results about Zumkeller numbers and suggested some conjectures in order to understand Zumkeller numbers.

In section 2 of this paper we shall study some properties of Zumkeller numbers. In section 3 we shall study the relations between practical numbers and Zumkeller numbers. We shall also settle a conjecture from [4] in this section. In section 4 we shall define and study the properties of half-Zumkeller numbers and prove results analogous to results about Zumkeller numbers for half-Zumkeller numbers. We shall also make substantial contributions regarding the second conjecture from [4].

2 Zumkeller numbers

Definition 1 *A positive integer n is said to be a Zumkeller number if the positive factors of n can be partitioned into two disjoint parts so that the sums of the two parts are equal. We shall call such a partition a Zumkeller partition.*

6, 12, 20, 24, 28, 30, 40 are the first few Zumkeller numbers. We shall start with a few simple facts. We shall also designate some of the known results as facts and for some of the facts we shall provide the proofs.

Let $\sigma(n)$ represent the sum of all positive factors of n .

Fact 1 *Let the prime factorization of n be $\prod_{i=1}^m p_i^{k_i}$. Then*

$$\sigma(n) = \prod_{i=1}^m \left(\sum_{j=0}^{k_i} p_i^j \right)$$

and

$$\frac{\sigma(n)}{n} = \prod_{i=1}^m \left(\sum_{j=0}^{k_i} p_i^{j-k_i} \right) = \prod_{i=1}^m \left(\sum_{j=0}^{k_i} p_i^{-j} \right) \leq \prod_{i=1}^m \frac{p_i}{p_i - 1}.$$

$\sigma(n)$ of a Zumkeller number n cannot be odd. This is observed in the next fact.

Fact 2 [1] *If n is a Zumkeller number, then $\sigma(n)$ must be even. Therefore the number of positive odd factors of n must be even.*

Proof. If $\sigma(n)$ is odd, then it is impossible to partition the positive factors of n into two equal-summed parts. If the number of positive odd factors of n is odd, then $\sigma(n)$ is odd. ■

The following fact gives a necessary and sufficient condition for n to be a Zumkeller number.

Fact 3 *n is a Zumkeller number if and only if $\frac{\sigma(n)-2n}{2}$ is either zero or is a sum of distinct positive factors of n excluding n itself.*

Proof. n is a Zumkeller number if and only if there exists a which is either 0 or a sum of some factors of n excluding n itself such that

$$n + a = \sigma(n) - (n + a). \quad (1)$$

This is equivalent to $\frac{\sigma(n)-2n}{2} = a$. ■

Let us state a simple necessary condition for a number to be Zumkeller. This says that every Zumkeller number is abundant.

Fact 4 [1] *If n is a Zumkeller number, then $\sigma(n) \geq 2n$.*

This follows from Fact 3.

Based on Fact 2, we shall now give a necessary condition for an even number to be Zumkeller.

Fact 5 *Let the prime factorization of an even Zumkeller number n be $2^k p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$. Then at least one of k_i must be odd.*

Proof. Note that the number of positive odd factors of n is $(k_1+1)(k_2+1) \cdots (k_m+1)$. At least one of k_i must be odd in order to make the number $(k_1+1)(k_2+1) \cdots (k_m+1)$ be even. ■

The following fact from [1] gives a method of generating new Zumkeller numbers from a known Zumkeller number.

Fact 6 [1] *If n is a Zumkeller number and p is a prime with $(n, p) = 1$, then np^l is Zumkeller for any positive integer l .*

Proof. Since n is a Zumkeller number, the set of all positive factors of n , denoted by D_0 can be partitioned into two disjoint parts A_0 and B_0 so that the sums of the two parts are equal. Group the factors of np^l into $l+1$ groups $D_0, D_1, \dots, D_l$ according to

how many factors of p they admit, i.e., D_i consists of all positive factors of np^l admitting i factors of p for every i , $0 \leq i \leq l$. Then each D_i can be partitioned into two disjoint parts so that the sum of these two parts equal according to the Zumkeller partition of $D_0 = A_0 \cup B_0$. ■

We give another method of generating new Zumkeller numbers from a known Zumkeller number.

Proposition 7 *Let n be a Zumkeller number and $p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$ be the prime factorization of n . Then for any positive integers $l_1, \dots, l_m$,*

$$p_1^{k_1+l_1(k_1+1)} p_2^{k_2+l_2(k_2+1)} \cdots p_m^{k_m+l_m(k_m+1)}$$

is Zumkeller.

Proof. It is sufficient to show that $p_1^{k_1+l_1(k_1+1)} p_2^{k_2} \cdots p_m^{k_m}$ is Zumkeller if $n = p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$ is Zumkeller. Assume that $n = p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$ is Zumkeller. Then the set of all positive factors of n , denoted by D_0 can be partitioned into two parts A_0 and B_0 so that the sums of these two parts are equal. Note that the factors of $p_1^{k_1+l_1(k_1+1)} p_2^{k_2} \cdots p_m^{k_m}$ can be partitioned into $l_1 + 1$ disjoint groups D_i , $0 \leq i \leq l_1$, where elements in D_i , $1 \leq i \leq l_1$ are obtained by multiplying elements in D_0 with $p_1^{i(k_1+1)}$. Note that every element in D_i admits at least $i(k_1 + 1)$ and at most $k_1 + i(k_1 + 1)$ factors of p_1 . So for $i \neq j$, D_i and D_j are disjoint. Since D_0 can be partitioned into two disjoint parts A_0 and B_0 so that the sums of the two parts are equal, all the other D_i 's can be partitioned into two disjoint parts A_i and B_i correspondingly so that the sums of these two parts equal. This proves that $p_1^{k_1+l_1(k_1+1)} p_2^{k_2} \cdots p_m^{k_m}$ is a Zumkeller number. ■

3 Practical numbers and Zumkeller numbers

Fact 3 states that if a certain number related to n is a sum of distinct positive factors of n , then n is Zumkeller. From the definition of practical numbers (given below), it is natural to consider the relations between practical numbers and Zumkeller numbers. We shall do this in this section. Practical numbers were introduced in [4].

Definition 2 *A positive integer n is said to be a practical number if all positive integers less than n can be represented as sums of distinct factors of n .*

As Stewart (1954) in [5] showed, it is straightforward to determine whether a number is practical from its prime factorization.

Fact 8 [5] *A positive integer n with the prime factorization $p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$ and $p_1 < p_2 < \cdots < p_m$ is a practical number if and only if $p_1 = 2$ and for $1 \leq i \leq m - 1$, $p_{i+1} \leq \sigma(p_1^{k_1} \cdots p_i^{k_i}) + 1$.*

The following result from [3] and [5] says that every positive integer $\leq n$ is a sum of distinct factors of n if and only if every positive integer $\leq \sigma(n)$ is a sum of distinct factors of n .

Proposition 9 [3] *A positive integer n is a practical number if and only if every positive integer $\leq \sigma(n)$ is a sum of distinct factors of n .*

Note that every practical number is even. Also note that for every positive integer k , 2^k is practical. The next result gives a sufficient condition for the product of 2^k and a prime $p \neq 2$ to be a Zumkeller number.

Fact 10 [1] *For any prime $p \neq 2$ and positive integer k with $p \leq 2^{k+1} - 1$, $2^k p$ is a Zumkeller number.*

With the aim of generalizing the above result for all practical numbers, the following problem was proposed as a conjecture in [1].

Conjecture 1 [1] *Let n be a practical number and p be a prime with $(n, p) = 1$, and $p \leq \sigma(n)$. Then, np is Zumkeller.*

Clark et al., suggested that some further restrictions on $\sigma(n)$ or $\frac{\sigma(n)}{n}$ are possibly needed for the conjecture to be true.

We shall prove a comprehensive theorem (Theorem 12) that settles this conjecture. We shall first find necessary and sufficient conditions for a practical number to be a Zumkeller number.

Proposition 11 *A practical number n is a Zumkeller number if and only if $\sigma(n)$ is even.*

Proof. If n is Zumkeller, then $\sigma(n)$ is even by Fact 2.

If $\sigma(n)$ is even, then $\frac{\sigma(n)}{2}$ is a positive integer smaller than $\sigma(n)$. Since n is practical, by Proposition 9, $\frac{\sigma(n)}{2}$ is the sum of some positive factors of n . Therefore, the set of all positive factors of n can be partitioned into two equal-summed parts and n is Zumkeller. ■

Theorem 12 *Let n be a practical number and p be a prime with $(n, p) = 1$. Let l be a positive integer. Then,*

(i) *If $\sigma(n)$ is even, then np^l is Zumkeller.*

(ii) *If $\sigma(n)$ is odd, then np^l is Zumkeller if and only if $p \leq \sigma(n)$ and l is odd.*

Proof. (i) By Proposition 11 and Fact 6, np^l is Zumkeller.

(ii) Let $\sigma(n)$ be odd. We first apply the conclusion in Case 1 to show that np^l is Zumkeller if $p \leq \sigma(n)$ and l is odd. Since n is practical and $p \leq \sigma(n)$, by Fact 8, np^l is practical. Since l is odd and p is odd, $\sum_{i=0}^l p^i$ is even and consequently, $\sigma(np^l) = (\sum_{i=0}^l p^i)\sigma(n)$ is even. By Proposition 11, we obtain that np^l is Zumkeller.

Now we shall show that if np^l is Zumkeller then, $p \leq \sigma(n)$ and l is odd. Let np^l be Zumkeller. Then for every integer i , $0 \leq i \leq l$, there exist a_i, b_i each of which is either 0 or a sum of some factors of n such that

$$\sum_{i=0}^l p^i(a_i - b_i) = 0;$$

and

$$a_i + b_i = \sigma(n).$$

Therefore, p divides $a_0 - b_0$. Since $a_0 + b_0 = \sigma(n)$ is odd, $a_0 - b_0 \neq 0$. Therefore, $|a_0 - b_0| \geq p$. Hence,

$$\sigma(n) = a_0 + b_0 \geq |a_0 - b_0| \geq p.$$

Since np^l is Zumkeller, $\sigma(np^l) = (\sum_{i=0}^l p^i)\sigma(n)$ must be even by Fact 2. Since $\sigma(n)$ is odd, $\sum_{i=0}^l p^i$ must be even. Therefore, l must be odd. ■

In Proposition 11 we showed that every practical number n with even $\sigma(n)$ is Zumkeller. But,

Remark 13 *There are Zumkeller numbers that are not practical numbers.*

All odd Zumkeller numbers are not practical numbers. For example,

$$\frac{\sigma(945) - 2 \cdot 945}{2} = \frac{(1 + 3 + 3^2 + 3^3) \cdot (1 + 5) \cdot (1 + 7) - 1890}{2} = 15$$

is a factor of 945. By Fact 3, 945 is a Zumkeller number. But 945 is not a practical number since all practical numbers are even.

Also, $70 = 2 \cdot 5 \cdot 7$ is an even Zumkeller number that is not practical. In fact,

$$\frac{\sigma(70) - 2 \cdot 70}{2} = \frac{(1 + 2) \cdot (1 + 5) \cdot (1 + 7) - 140}{2} = \frac{144 - 140}{2} = 2$$

is a factor of 70. By Fact 3, 70 is a Zumkeller number. But it is not a practical number by Fact 8.

Now we shall develop several results for Zumkeller numbers and these will be used later. A finer analysis of the proof of (ii) in the above theorem, gives us the following result.

Proposition 14 *Let l be a positive integer. Let n be a non-Zumkeller number and p be a prime with $(n, p) = 1$. If np^l is Zumkeller, then $p \leq \sigma(n)$. If np^l is Zumkeller and $\sigma(n)$ is odd, then l is odd.*

Proof. This is similar to the proof of the necessity part of Theorem 12 (ii). The only modification is to replace ‘Since $a_0 + b_0 = \sigma(n)$ is odd, $a_0 - b_0 \neq 0$ ’ by ‘Since $a_0 + b_0 = \sigma(n)$ and n is not Zumkeller, $a_0 - b_0 \neq 0$ ’. ■

We will now give several necessary and sufficient conditions for np to be a Zumkeller number.

Proposition 15 *Let n be a positive integer and p be a prime with $(n, p) = 1$. Then the following conditions are equivalent:*

- (i) np is Zumkeller.
- (ii) The set of all positive factors of n can be partitioned into two disjoint parts D_1 and D_2 such that $p(\sum_{d \in D_2} d - \sum_{d \in D_1} d)$ can be written as a sum of some factors of n minus the sum of the rest of the factors of n .
- (iii) The set of all positive factors of n can be partitioned into two disjoint parts D_1 and D_2 such that $\frac{(p+1)(\sum_{d \in D_2} d - \sum_{d \in D_1} d)}{2}$ can be written as a sum of some elements in D_2 minus a sum of some elements in D_1 .
- (iv) The set of all positive factors of n can be partitioned into 4 disjoint parts A_1, A_2, A_3 and A_4 such that $(p+1) \sum_{d \in A_1} d + (p-1) \sum_{d \in A_2} d = (p+1) \sum_{d \in A_3} d + (p-1) \sum_{d \in A_4} d$.

Proof. np is Zumkeller if and only if the set D_0 of all positive factors of n can be partitioned into $D_0 = D_1 \cup D_2$ and $D_0 = D_3 \cup D_4$ such that

$$p\left(\sum_{d \in D_1} d\right) + \sum_{d \in D_4} d = p\left(\sum_{d \in D_2} d\right) + \sum_{d \in D_3} d.$$

This can be rewritten as

$$p\left(\sum_{d \in D_2} d - \sum_{d \in D_1} d\right) + \left(\sum_{d \in D_3} d - \sum_{d \in D_4} d\right) = 0.$$

This is equivalent to

$$p\left(\sum_{d \in D_2} d - \sum_{d \in D_1} d\right) = -\left(\sum_{d \in D_3} d - \sum_{d \in D_4} d\right). \quad (2)$$

This is equivalent to (ii).

Now, by adding $\sum_{d \in D_2} d - \sum_{d \in D_1} d$ to both sides of the above equation we get,

$$\begin{aligned} p\left(\sum_{d \in D_2} d - \sum_{d \in D_1} d\right) + \left(\sum_{d \in D_2} d - \sum_{d \in D_1} d\right) &= \left(\sum_{d \in D_2} d - \sum_{d \in D_1} d\right) - \left(\sum_{d \in D_3} d - \sum_{d \in D_4} d\right) \\ &= \left(\sum_{d \in D_2} d - \sum_{d \in D_3} d\right) - \left(\sum_{d \in D_1} d - \sum_{d \in D_4} d\right) \\ &= \left(\sum_{d \in D_2} d - \sum_{d \in D_3 \cap D_2} d\right) - \sum_{d \in D_3 \cap D_1} d \\ &\quad - \left(\sum_{d \in D_1} d - \sum_{d \in D_4 \cap D_1} d\right) + \sum_{d \in D_4 \cap D_2} d \\ &= \sum_{d \in D_4 \cap D_2} d - \sum_{d \in D_3 \cap D_1} d - \sum_{d \in D_3 \cap D_1} d + \sum_{d \in D_4 \cap D_2} d \\ &= 2\left(\sum_{d \in D_4 \cap D_2} d - \sum_{d \in D_3 \cap D_1} d\right). \end{aligned}$$

Thus,

$$\frac{(p+1)\left(\sum_{d \in D_2} d - \sum_{d \in D_1} d\right)}{2} = \sum_{d \in D_4 \cap D_2} d - \sum_{d \in D_3 \cap D_1} d.$$

This is equivalent to (iii). To go in the converse direction one needs to take $D_3 = (D_3 \cap D_1) \cup (D_2 - D_4 \cap D_2)$ and $D_4 = (D_4 \cap D_2) \cup (D_1 - D_3 \cap D_1)$.

By adding $p(\sum_{d \in D_3} d - \sum_{d \in D_4} d)$ to both sides of equation (2) we get that equation (2) is equivalent to

$$p\left(\sum_{d \in D_2} d - \sum_{d \in D_1} d\right) + p\left(\sum_{d \in D_3} d - \sum_{d \in D_4} d\right) = -\left(\sum_{d \in D_3} d - \sum_{d \in D_4} d\right) + p\left(\sum_{d \in D_3} d - \sum_{d \in D_4} d\right).$$

This is equivalent to

$$p\left(\sum_{d \in D_2} d - \sum_{d \in D_4} d + \sum_{d \in D_3} d - \sum_{d \in D_1} d\right) = (p-1)\left(\sum_{d \in D_3} d - \sum_{d \in D_4} d\right).$$

This is equivalent to

$$p\left(\sum_{d \in D_2} d - \sum_{d \in D_2 \cap D_4} d - \sum_{d \in D_1 \cap D_4} d + \sum_{d \in D_3} d - \sum_{d \in D_1 \cap D_3} d - \sum_{d \in D_1 \cap D_4} d\right) = (p-1)\left(\sum_{d \in D_3} d - \sum_{d \in D_4} d\right).$$

This is equivalent to

$$p\left(\sum_{d \in D_2 \cap D_3} d - \sum_{d \in D_1 \cap D_4} d + \sum_{d \in D_2 \cap D_3} d - \sum_{d \in D_1 \cap D_4} d\right) = (p-1)\left(\sum_{d \in D_3} d - \sum_{d \in D_4} d\right).$$

This is equivalent to

$$2p\left(\sum_{d \in D_2 \cap D_3} d - \sum_{d \in D_1 \cap D_4} d\right) = (p-1)\left(\sum_{d \in D_3} d - \sum_{d \in D_4} d\right).$$

This is equivalent to

$$\begin{aligned} (p+1)\left(\sum_{d \in D_2 \cap D_3} d - \sum_{d \in D_1 \cap D_4} d\right) &= (p-1)\left(\sum_{d \in D_3} d - \sum_{d \in D_4} d - \sum_{d \in D_2 \cap D_3} d + \sum_{d \in D_1 \cap D_4} d\right) \\ &= (p-1)\left(\sum_{d \in D_1 \cap D_3} d - \sum_{d \in D_2 \cap D_4} d\right). \end{aligned}$$

This is equivalent to

$$(p+1)\sum_{d \in D_2 \cap D_3} d + (p-1)\sum_{d \in D_2 \cap D_4} d = (p+1)\sum_{d \in D_1 \cap D_4} d + (p-1)\sum_{d \in D_1 \cap D_3} d.$$

By letting $A_1 = D_2 \cap D_3$, $A_2 = D_2 \cap D_4$, $A_3 = D_1 \cap D_4$, and $A_4 = D_1 \cap D_3$ and observing that $\{A_1, A_2, A_3, A_4\}$ is a partition of the positive factors of n we see that the above equation is equivalent to (iv). ■

Remark 16 *If the set of all positive factors of n can be partitioned into two disjoint parts D_1 and D_2 such that $\sum_{d \in D_2} d - \sum_{d \in D_1} d$ is small, then it is usually easy to check condition (iii) in Proposition 15.*

Let us see an application of Proposition 15.

Fact 17 $2 \times 5^2 \times 7^2 \times p$ is Zumkeller for $p = 11, 13$.

Proof. The set of all positive factors of $2 \times 5^2 \times 7^2$ can be partitioned into two disjoint parts $D_1 = \{2450, 98, 50, 35, 10, 5, 2\}$ and $D_2 = \{1225, 490, 350, 245, 175, 70, 49, 25, 14, 7, 1\}$ and $\sum_{d \in D_2} d - \sum_{d \in D_1} d = 1$.

$$\frac{(11+1)(\sum_{d \in D_2} d - \sum_{d \in D_1} d)}{2} = 6 = (7+1) - 2$$
is a sum of some elements in D_2 minus a sum of some elements in D_1 and it satisfies Proposition 15(iii) for $p = 11$. So $2 \times 5^2 \times 7^2 \times 11$ is Zumkeller.

$\frac{(13+1)(\sum_{d \in D_2} d - \sum_{d \in D_1} d)}{2} = 7$ is a sum of some elements in D_2 and it satisfies Proposition 15(iii) for $p = 13$. So $2 \times 5^2 \times 7^2 \times 13$ is Zumkeller. ■

One can use the above method to show that Fact 17 holds for more prime numbers.

In Proposition 1.4 of [3] a condition on the existence of factors with certain nice properties was shown to be sufficient for an integer to be a practical number. We shall prove a similar theorem for Zumkeller numbers. In this case we shall also develop a method of finding a Zumkeller partition.

Proposition 18 *If $1 = a_1 < a_2 < \dots < a_k = n$ are all factors of n with $a_{i+1} \leq 2a_i$ for all i and $\sigma(n)$ is even, then n is Zumkeller.*

Proof. Starting with a positive sign for $a_k = n$, we will assign positive or negative signs to each a_i and show that the sum of all a_i , $1 \leq i \leq k$ with the positive or negative signs assigned is 0. Then it will imply that $\sigma(n)$ can be partitioned into two equal-summed parts.

a_k is assigned positive sign, and a_{k-1} is assigned the negative sign. Note that $0 < a_k - a_{k-1} \leq a_{k-1}$ since $a_k \leq 2a_{k-1}$. Since the current sum $a_k - a_{k-1}$ is positive, we assign the negative sign to a_{k-2} . Then $-a_{k-2} < a_k - a_{k-1} - a_{k-2} \leq a_{k-1} - a_{k-2} \leq a_{k-2}$ since $a_{k-1} \leq 2a_{k-2}$. If $a_k - a_{k-1} - a_{k-2} \geq 0$, we assign the negative sign to a_{k-3} . Otherwise we assign the positive sign to a_{k-3} . Let s_i be the current sum up to a_i . In general, the sign assigned to a_{i-1} is opposite of the sign of s_i . Let us show inductively that $|s_i| \leq a_i$ for $1 \leq i \leq k$. It is true for $i = k$. Assume that $|s_{i+1}| \leq a_{i+1}$. Since the sign of a_i is opposite of the sign of s_{i+1} , $|s_i| = ||s_{i+1}| - a_i|$. Note that $-a_i \leq |s_{i+1}| - a_i \leq a_{i+1} - a_i \leq a_i$ since $a_{i+1} \leq 2a_i$. Therefore $|s_i| \leq a_i$. So $|s_1| \leq a_1 = 1$. Since $\sigma(n)$ is even, s_1 , which is obtained by assigning a positive or negative sign to each of the terms in $\sigma(n)$ is even as well. So $s_1 = 0$. This implies that $\sigma(n)$ can be partitioned into two equal-summed parts, i.e., n is Zumkeller. ■

Clearly, if n is an integer for which we can find factors $1 = b_1 < b_2 < \dots < b_k = n$ with the property that $b_{i+1} \leq 2b_i$ for all i and $\sigma(n)$ is even, then the hypothesis of Proposition 18 is satisfied (for all factors of n). Hence such an n is Zumkeller.

Remark 19 *In [3], $n!$ was shown to be a practical number for all $n \geq 3$. We shall now apply Proposition 18 and show that $n!$ is a Zumkeller number for all $n \geq 3$.*

Proof. If $1 = a_1 < a_2 < \dots < a_k = n!$ are all factors of $n!$, then clearly $a_{i+1} \leq 2a_i$ for all i (Note that $1, 2, 3, \dots, n, 2n, 3n, \dots, (n-1)n, 2(n-1)n, 3(n-1)n, \dots, (n-2)(n-$

1) n , $2(n-2)(n-1)n$, $3(n-2)(n-1)n$, $\dots$, $\prod_{i=3}^n i$, $2 \prod_{i=3}^n i = n!$ are some of the factors of $n!$.) By Proposition 18, it is sufficient to show that $\sigma(n!)$ is even for $n \geq 3$. Let p be the largest prime $\leq n$. If $n! = p_1^{k_1} p_2^{k_2} \cdots p_{m_n}^{k_{m_n}}$ is the prime factorization of $n!$ with $2 = p_1 < p_2 < \cdots < p_m$, then clearly $p_m = p$. If $k_m \geq 2$, then $2p \leq n$ and by Bertrand's postulate, there exists a prime number q such that $\frac{n}{2} < q \leq n$ and this contradicts the choice of p . So $k_m = 1$. This implies that $\sigma(n!)$ is even. ■

We shall now discuss odd Zumkeller numbers. As stated in [1], odd Zumkeller numbers do exist. In fact, the first several odd abundant numbers with even σ -value are Zumkeller. Using Proposition 7, starting with an odd Zumkeller number one can produce an infinite sequence of Zumkeller numbers. The next result describes the prime factors of an odd Zumkeller number with a small number of prime factors.

Fact 20 *Let the prime factorization of an odd number n be $p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$, where $3 \leq p_1 < p_2 < \cdots < p_m$. If n is Zumkeller, then*

$$\prod_{i=1}^m \frac{p_i}{p_i - 1} \geq 2.$$

and m is at least 3. In particular:

1. If $m \leq 6$, then $p_1 = 3$, $p_2 = 5, 7$ or 11 .
2. If $m \leq 4$, then $p_1 = 3$, $p_2 = 5$.
3. If $m = 3$, then $p_1 = 3$, $p_2 = 5$, and $p_3 = 7$ or 11 or 13 .

Proof. If n is Zumkeller, then by Facts 4 and 1,

$$2p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m} = 2n \leq \sigma(n) = \prod_{i=1}^m \left(\sum_{j=0}^{k_i} p_i^j \right).$$

Dividing both sides by $p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$, we get

$$2 \leq \prod_{i=1}^m \left(\sum_{j=0}^{k_i} p_i^{j-k_i} \right) \leq \prod_{i=1}^m \frac{p_i}{p_i - 1}.$$

If $m \leq 2$, then

$$\prod_{i=1}^m \frac{p_i}{p_i - 1} \leq \frac{3}{2} \times \frac{5}{4} < 2.$$

So $m \geq 3$. The truth of 1 – 3 follows by verifying the condition $\prod_{i=1}^m \frac{p_i}{p_i - 1} \geq 2$ directly as given below.

1. Let $m \leq 6$. If $p_1 \neq 3$, then $p_1 \geq 5$ and

$$\prod_{i=1}^m \frac{p_i}{p_i - 1} \leq \frac{5}{4} \times \frac{7}{6} \times \frac{11}{10} \times \frac{13}{12} \times \frac{17}{16} \times \frac{19}{18} < 2.$$

Therefore, $p_1 = 3$. If $p_2 > 11$, then $p_2 \geq 13$ and

$$\prod_{i=1}^m \frac{p_i}{p_i - 1} \leq \frac{3}{2} \times \frac{13}{12} \times \frac{17}{16} \times \frac{19}{18} \times \frac{23}{22} \times \frac{29}{28} < 2.$$

Hence, $p_2 \leq 11$. This implies that $p_2 = 5, 7$ or 11 .

2. Let $m \leq 4$. By 1, $p_1 = 3$. If $p_2 > 7$, then $p_2 \geq 11$, so

$$\prod_{i=1}^m \frac{p_i}{p_i - 1} \leq \frac{3}{2} \times \frac{11}{10} \times \frac{13}{12} \times \frac{17}{16} < 2.$$

Therefore, $p_2 \leq 7$. This implies that $p_2 = 5$ or 7 .

3. Let $m = 3$. By 1, $p_1 = 3$. If $p_2 \neq 5$, then $p_2 \geq 7$ and $p_3 \geq 11$. So

$$\prod_{i=1}^3 \frac{p_i}{p_i - 1} \leq \frac{3}{2} \times \frac{7}{6} \times \frac{11}{10} < 2.$$

Hence, $p_2 = 5$.

If $p_3 \geq 17$, then

$$\prod_{i=1}^3 \frac{p_i}{p_i - 1} \leq \frac{3}{2} \times \frac{5}{4} \times \frac{17}{16} < 2.$$

Hence, $p_3 < 17$ and consequently $p_3 = 7, 11$ or 13 . ■

We do not know too much more about odd Zumkeller numbers.

4 Half-Zumkeller numbers

Definition 3 A positive integer n is said to be half-Zumkeller number if the proper positive factors of n can be partitioned into two disjoint parts so that the sums of the two parts are equal.

We shall start with some simple observations.

Fact 21 A positive integer n is half-Zumkeller if and only if $\frac{\sigma(n)-n}{2}$ is the sum of some distinct positive proper factors of n .

Fact 22 A positive even integer n is half-Zumkeller if and only if $\frac{\sigma(n)-2n}{2}$ is zero or the sum of some distinct factors of n excluding n and $\frac{n}{2}$.

Proof. Let n be even. n is half-Zumkeller if and only if there exists a which is zero or the sum of some factors of n excluding n and $\frac{n}{2}$ such that

$$\frac{n}{2} + a = \sigma(n) - n - \left(\frac{n}{2} + a\right).$$

Therefore, $a = \frac{\sigma(n) - 2n}{2}$. ■

Fact 23 $2 \cdot 5 \cdot 7$, $2 \cdot 5^2 \cdot 7$, and $2 \cdot 5 \cdot 7^2$ are half-Zumkeller numbers.

Proof. $2 \cdot 5 \cdot 7$ is half-Zumkeller since

$$\frac{\sigma(2 \cdot 5 \cdot 7) - 2(2 \cdot 5 \cdot 7)}{2} = 2$$

is a sum of factors of $2 \cdot 5 \cdot 7$ excluding $2 \cdot 5 \cdot 7$ and $5 \cdot 7$ (by Fact 22). Similarly, $2 \cdot 5^2 \cdot 7$ is half-Zumkeller since

$$\frac{\sigma(2 \cdot 5^2 \cdot 7) - 2(2 \cdot 5^2 \cdot 7)}{2} = 22 = 2 \cdot 7 + 7 + 1$$

is a sum of factors of $2 \cdot 5^2 \cdot 7$ excluding $2 \cdot 5^2 \cdot 7$ and $5^2 \cdot 7$; and $2 \cdot 5 \cdot 7^2$ is half-Zumkeller since

$$\frac{\sigma(2 \cdot 5 \cdot 7^2) - 2(2 \cdot 5 \cdot 7^2)}{2} = 23 = 2 \cdot 7 + 7 + 2$$

is a sum of factors of $2 \cdot 5 \cdot 7^2$ excluding $2 \cdot 5 \cdot 7^2$ and $5 \cdot 7^2$. ■

The next four results are pointed out in [1].

Fact 24 [1] *If n is odd and half-Zumkeller, then n is a perfect square.*

Proof. If n is odd and half-Zumkeller, then $\sigma(n) - n$ must be even and $\sigma(n)$ must be odd. Let the prime factorization of n be $\prod_{i=1}^m p_i^{k_i}$. Then $\sigma(n) = \prod_{i=1}^m (\sum_{j=1}^{k_i} p_i^j)$. If $\sigma(n)$ is odd, then all k_i must be even. So n is a perfect square. ■

Fact 25 [1] *If m and n are half-Zumkeller numbers with $(m, n) = 1$, then mn is half-Zumkeller.*

Fact 26 [1] *If n is even and half-Zumkeller, then n is Zumkeller.*

Proof. Let D be the set of all positive factors of n . If n is even and half-Zumkeller, then there exists $A \subset D \setminus \{n, \frac{n}{2}\}$ such that

$$\frac{n}{2} + \sum_{a \in A} a = \sum_{b \in D, b \notin \{n, \frac{n}{2}\} \cup A} b.$$

Adding $\frac{n}{2}$ to both sides, we have

$$n + \sum_{a \in A} a = \frac{n}{2} + \sum_{b \in D, b \notin \{n, \frac{n}{2}\} \cup A} b.$$

This means that all positive factors of n are partitioned into two equal-summed parts and n is Zumkeller. ■

Remark 27 [1] *Let n be even. n is half-Zumkeller if and only if n admits a Zumkeller partition such that n and $\frac{n}{2}$ are in distinct parts.*

Proof. Let n be even. Let D be the set of all positive factors of n . n is half-Zumkeller if and only if there exists $A \subset D \setminus \{n, \frac{n}{2}\}$ such that

$$\frac{n}{2} + \sum_{a \in A} a = \sum_{b \in D, b \notin \{n, \frac{n}{2}\} \cup A} b.$$

That is.,

$$n + \sum_{a \in A} a = \frac{n}{2} + \sum_{b \in D, b \notin \{n, \frac{n}{2}\} \cup A} b.$$

This is equivalent to saying that n admits a Zumkeller partition such that n and $\frac{n}{2}$ are in distinct parts. ■

The following conjecture is proposed in [1].

Conjecture 2 [1] *If n is even and Zumkeller, then n is half-Zumkeller.*

In two of the next three results we shall verify that the conjecture is true in some cases.

Proposition 28 *Let n be even and Zumkeller. If $\sigma(n) < 3n$, then n is half-Zumkeller.*

Proof. If n is Zumkeller, by Fact 3, $\frac{\sigma(n)-2n}{2}$ is either zero or the sum of some factors of n excluding n . If $\sigma(n) < 3n$, then $\frac{\sigma(n)-2n}{2} < \frac{n}{2}$. So $\frac{\sigma(n)-2n}{2}$ excludes $\frac{n}{2}$ as well. Then $\frac{\sigma(n)-2n}{2}$ is zero or the sum of some factors of n excluding n and $\frac{n}{2}$. By Fact 22, n is half-Zumkeller. ■

Proposition 29 *Let n be even. Then, n is Zumkeller if and only if n is either half-Zumkeller or $\frac{\sigma(n)-3n}{2}$ is 0 or is a sum of some factors of n excluding n and $\frac{n}{2}$.*

Proof. Let n be even. If n is Zumkeller but not half-Zumkeller, then by Remark 27, any Zumkeller partition of the positive factors of n must have n and $\frac{n}{2}$ in the same parts. In other words, there exists a which is either 0 or a sum of some factors of n excluding n and $\frac{n}{2}$ such that

$$2(n + \frac{n}{2} + a) = \sigma(n).$$

So, $a = \frac{\sigma(n)-3n}{2}$. Therefore, $\frac{\sigma(n)-3n}{2}$ is zero or a sum of some factors of n excluding n and $\frac{n}{2}$.

If n is half-Zumkeller, then n is Zumkeller by Fact 26. If $\frac{\sigma(n)-3n}{2}$ is zero or a sum of some factors of n excluding n and $\frac{n}{2}$, then

$$\frac{\sigma(n) - 2n}{2} = \frac{\sigma(n) - 3n}{2} + \frac{n}{2}$$

is a sum of some factors of n excluding n . By Fact 3, n is Zumkeller. ■

Proposition 30 *If 2 divides n , 3 divides n , n is Zumkeller, and $\sigma(n) < \frac{10n}{3}$, then n is half-Zumkeller.*

Proof. If n is not half-Zumkeller, by Proposition 29, $(\sigma(n) - 3n)/2$ is 0 or a sum of some factors of n excluding n and $n/2$. Then,

$$(\sigma(n) - 2n)/2 = (\sigma(n) - 3n)/2 + n/3 + n/6.$$

Since $\sigma(n)/n < 10/3$ we have that $(\sigma(n) - 3n)/2 < n/6$. Hence $(\sigma(n) - 2n)/2$ is a sum of some factors of n excluding n and $n/2$. By Fact 22, n is half Zumkeller. This is a contradiction. ■

The next proposition identifies some half-Zumkeller numbers.

Proposition 31 *If n is Zumkeller, then $2n$ is half-Zumkeller.*

Proof. Let $n = 2^k L$ with k a nonnegative integer and L an odd number, be a Zumkeller number. Then all positive factors of n can be partitioned into two disjoint equal-summed parts D_1 and D_2 . Observe that every factor of $2n$ which is not a factor of n can be written as $2^{k+1}l$ where l is a factor of L . Observe that $2^k l$ is either in D_1 or D_2 . Without loss of generality, assume that $2^k l$ is in D_1 . In this case, we move $2^k l$ to D_2 and add $2^{k+1}l$ to D_1 . Perform this procedure to all factors of $2n$ which are not factors of n except $2n$ itself. This procedure will yield an equal-summed partition of all factors of $2n$ except $2n$ itself. This shows that $2n$ is half-Zumkeller. ■

The following remark is an immediate consequence of the above proposition. It identifies the prime factorization of an even Zumkeller number that is not half-Zumkeller.

Remark 32 *Let n be even and the prime factorization of n be $2^k p_1^{k_1} \cdots p_m^{k_m}$. If n is Zumkeller but not half-Zumkeller, then $2^i p_1^{k_1} \cdots p_m^{k_m}$ is not Zumkeller for any $i \leq k - 1$, and $2^i p_1^{k_1} \cdots p_m^{k_m}$ is half-Zumkeller for any $i \geq k + 1$. If Conjecture 2 is not true, then a counterexample to the Conjecture must have a prime factorization of $2^k p_1^{k_1} \cdots p_m^{k_m}$ such that k is the minimum integer i such that $2^i p_1^{k_1} \cdots p_m^{k_m}$ is Zumkeller.*

In order to investigate whether there are even Zumkeller numbers that are not half-Zumkeller, we shall see if all the results that are true for Zumkeller numbers are true for even half-Zumkeller numbers also.

Results analogous to Propositions 6 and 7, are true for half-Zumkeller numbers also.

Proposition 33 *Let n be an even half-Zumkeller number and p be a prime with $(n, p) = 1$. Then np^l is half-Zumkeller for any positive integer l .*

Proof. Since n is an even half-Zumkeller number, the set of all positive factors of n , denoted by D_0 can be partitioned into two disjoint parts A_0 and B_0 so that the sums of the two parts are equal and n and $\frac{n}{2}$ are in distinct parts (by Remark 27).

Group the factors of np^l into $l + 1$ groups $D_0, D_1, \dots, D_l$ according to how many factors of p they admit, i.e., D_i consists of all factors of np^l admitting i factors of p . Then each D_i can be partitioned into two disjoint parts so that the sums of the two parts are equal and np^i and $\frac{np^i}{2}$ are in distinct parts according to the Zumkeller partition of the set D_0 . Therefore all positive factors of np^l can be partitioned into two disjoint parts so that the sum of these two parts equal and np^l and $\frac{np^l}{2}$ are in distinct parts. By Remark 27, np^l is half-Zumkeller. ■

The following is a direct Corollary of Proposition 33.

Corollary 34 *If n is an even half-Zumkeller number and m is a positive integer with $(n, m) = 1$, then nm is half-Zumkeller.*

But this result is not true for odd half-Zumkeller numbers.

Remark 35 *If n is an odd half-Zumkeller number and m is a positive integer with $(m, n) = 1$, then mn need not be a half-Zumkeller number. For example, $3^2 \times 5^2$ is half-Zumkeller since all its proper positive factors can be partitioned into two equal parts: $75 + 9 + 5 = 45 + 25 + 15 + 3 + 1$. But $3^2 \times 5^2 \times 7$ is not half-Zumkeller by Fact 24.*

We shall prove Proposition 7 for even half-Zumkeller numbers.

Proposition 36 *Let n be an even half-Zumkeller number and the prime factorization of n be $p_1^{k_1} p_2^{k_2} \dots p_m^{k_m}$. Then for positive integers $l_1, \dots, l_m$,*

$$p_1^{k_1+l_1(k_1+1)} p_2^{k_2+l_2(k_2+1)} \dots p_m^{k_m+l_m(k_m+1)}$$

is half-Zumkeller.

Proof. It is sufficient to show that $p_1^{k_1+l_1(k_1+1)} p_2^{k_2} \dots p_m^{k_m}$ is half-Zumkeller if $n = p_1^{k_1} p_2^{k_2} \dots p_m^{k_m}$ is an even half-Zumkeller number. Assume that $n = p_1^{k_1} p_2^{k_2} \dots p_m^{k_m}$ is even and half-Zumkeller, then the set of all positive factors of n , denoted by D_0 can be partitioned into two disjoint parts A_0 and B_0 so that the sums of the two parts are equal and n and $\frac{n}{2}$ are in distinct parts (by Remark 27). Note that the factors of $p_1^{k_1+l_1(k_1+1)} p_2^{k_2} \dots p_m^{k_m}$ can be partitioned into $l_1 + 1$ disjoint groups D_i , $0 \leq i \leq l_1$, where elements in D_i are obtained by multiplying $p_1^{i(k_1+1)}$ with elements in D_0 . Using the partition A_0, B_0 of D_0 we can partition every D_i into two disjoint parts A_i and B_i so that the sums of the corresponding parts are equal and $np_1^{i(k_1+1)}$ and $\frac{np_1^{i(k_1+1)}}{2}$ are in distinct parts. Therefore, the set of all positive factors of $p_1^{k_1+l_1(k_1+1)} p_2^{k_2} \dots p_m^{k_m}$ can be partitioned into two disjoint equal-summed parts and $p_1^{k_1+l_1(k_1+1)} p_2^{k_2} \dots p_m^{k_m}$ and $\frac{p_1^{k_1+l_1(k_1+1)} p_2^{k_2} \dots p_m^{k_m}}{2}$ are in distinct parts. By Remark 27, $p_1^{k_1+l_1(k_1+1)} p_2^{k_2} \dots p_m^{k_m}$ is half-Zumkeller. ■

We shall now proceed to prove Proposition 11 and Theorem 12 for half-Zumkeller numbers.

We need a definition. This is similar to the concept of a practical number, but is related to half-Zumkeller numbers as practical numbers are related to Zumkeller numbers.

Definition 4 *We say that a positive integer n is a quasi-practical number if every positive integer $\leq \sigma(n) - n$ can be written as a sum of distinct positive factors of n excluding n .*

Clearly every prime number is a quasi-practical number. Some simple quasi-practical numbers are given by the following proposition.

Proposition 37 *For every nonnegative integer k , 2^k is quasi-practical.*

Proof. Since every integer $m < \sigma(2^k) - 2^k = 2^k - 1$ can be written as a sum of factors of 2^{k-1} (take the binomial expansion of m), 2^k is quasi-practical. ■

By Fact 21, if n is quasi-practical and $\sigma(n) - n$ is even, then n is half-Zumkeller.

Our next aim is to show that every practical number is quasi-practical. In order to show this we need a technical lemma.

Lemma 38 *Let p be a prime. Let $l, A_0, A_1, \dots, A_l$ be nonnegative integers. Every non-negative $M \leq A_0 + A_1p + \dots + A_lp^l$ can be written as $C_0 + C_1p + \dots + C_lp^l$ for some $C_0 \leq A_0, C_1 \leq A_1, \dots, C_l \leq A_l$ if and only if the following l conditions*

$$A_0 + 1 \geq p,$$

$$A_0 + A_1p + 1 \geq p^2,$$

.....

$$A_0 + A_1p + \dots + A_{l-1}p^{l-1} + 1 \geq p^l$$

are satisfied.

Proof. Necessity: If every nonnegative $M \leq A_0 + A_1p + \dots + A_lp^l$ can be written as $C_0 + C_1p + \dots + C_lp^l$ for some $C_0 \leq A_0, C_1 \leq A_1, \dots, C_l \leq A_l$, then for any $0 \leq i \leq l-1, A_0 + A_1p + \dots + A_ip^i + 1 \leq A_0 + A_1p + \dots + A_lp^l$ and this can be written as $C_0 + C_1p + \dots + C_lp^l$ for some $C_0 \leq A_0, C_1 \leq A_1, \dots, C_l \leq A_l$. Note that for some $i+1 \leq j \leq l, C_j \neq 0$ since, otherwise $C_0 + C_1p + \dots + C_ip^i < A_0 + A_1p + \dots + A_ip^i + 1$. Therefore, $A_0 + A_1p + \dots + A_ip^i + 1 \geq p^{j+1} \geq p^{i+1}$.

We thus have the necessity.

Sufficiency: We shall prove this by induction. For $l = 0$ there is nothing to be shown.

Let $l \geq 1$. Take a nonnegative integer $M \leq A_0 + A_1p + \dots + A_lp^l$.

Find the largest $C_l \leq A_l$ such that $C_lp^l \leq M$. If $C_l = A_l$, then, $M - A_lp^l \leq A_0 + A_1p + A_2p^2 + \dots + A_{l-1}p^{l-1}$ and the induction will take care of the rest. If $C_l < A_l$, then, $C_lp^l \leq M < (C_l + 1)p^l$. This implies that $M - C_lp^l < p^l$. But, from the hypothesis, $p^l - 1 \leq A_0 + A_1p + \dots + A_{l-1}p^{l-1}$. Again the induction will take care of the rest. ■

We are now ready to prove a result about quasi-practical numbers that will help us show that every practical number is quasi-practical.

Proposition 39 *Let n be a practical and quasi-practical number. Let p be a prime with $(n, p) = 1$ and l be a positive integer. Then np^l is quasi-practical if and only if $p \leq \sigma(n) + 1$.*

Proof. Necessity: If np^l is quasi-practical, then the positive integer $p - 1 < \sigma(np^l) - np^l$ is a sum of factors of np^l excluding np^l . Since $p - 1 < p$, $p - 1$ must be a sum of factors of n , then $p - 1 \leq \sigma(n)$. So $p \leq \sigma(n) + 1$.

Sufficiency: Let M be a positive integer less than $\sigma(np^l) - np^l$. Then

$$\begin{aligned} M < \sigma(np^l) - np^l &= \sigma(n) \left(\sum_{i=0}^l p^i \right) - np^l \\ &= \left(\sum_{i=0}^{l-1} \sigma(n)p^i \right) + (\sigma(n) - n)p^l. \end{aligned} \quad (3)$$

Since $\sigma(n) + 1 \geq p$, we get that

$$\sigma(n) + \sigma(n)p + 1 \geq p^2,$$

.....

$$\sigma(n) + \sigma(n)p + \dots + \sigma(n)p^{l-1} + 1 \geq p^l$$

Since n is practical every positive integer $\leq \sigma(n)$ is a sum of factors of n . Since n is also quasi-practical every positive integer $\leq \sigma(n) - n$ is a sum of factors of n excluding n . Applying Lemma 38 to (3), M can be written as $C_0 + C_1p + \dots + C_l p^l$ for some $C_0 \leq \sigma(n)$, $C_1 \leq \sigma(n)$, ..., $C_{l-1} \leq \sigma(n)$, $C_l \leq \sigma(n) - n$ if $p \leq \sigma(n) + 1$. Since n is a practical number and a quasi-practical number, then C_l is a sum of factors of n excluding n itself and each C_i , $1 \leq i \leq l - 1$ is a sum of factors of n . Therefore, M is a sum of factors of np^l excluding np^l itself. So np^l is quasi-practical if $p \leq \sigma(n) + 1$. ■

Now, we are ready to derive some consequences of the above result.

Proposition 40 *Every practical number is quasi-practical.*

Proof: By Fact 8 any practical number n looks like $n = p_1^{k_1} p_2^{k_2} \dots p_m^{k_m}$ and $p_1 < p_2 < \dots < p_m$ where $p_1 = 2$ and for $1 \leq i \leq m - 1$, $p_{i+1} \leq \sigma(p_1^{k_1} \dots p_i^{k_i}) + 1$. But 2^{k_1} is both practical and quasi-practical. Repeated use of Proposition 39 shows that n is also quasi-practical. ■

Remark 41 *A number is quasi-practical if and only if either it is a practical number or a prime number. In fact even quasi-practical numbers are same as practical numbers. The difficult part of this is Proposition 40. We omit the rest of the details.*

We shall now find necessary and sufficient conditions for a practical number to be a half-Zumkeller number.

Proposition 42 *A practical number n is a half-Zumkeller number if and only if $\sigma(n)$ is even.*

Proof. If n is practical and half-Zumkeller, it is an even half-Zumkeller number. So, n is Zumkeller by Fact 26 and $\sigma(n)$ is even by Fact 2.

If $\sigma(n)$ is even, then $\frac{\sigma(n)-n}{2}$ is an integer smaller than $\sigma(n) - n$. Since n is a practical number, it is quasi-practical by Proposition 40. Therefore $\frac{\sigma(n)-n}{2}$ is a sum of factors of n excluding n . By Fact 21, n is half-Zumkeller.

Theorem 43 *Let n be a practical number, l be a positive integer and p be a prime with $(n, p) = 1$.*

- (i). *If $\sigma(n)$ is even, then np^l is half-Zumkeller.*
- (ii). *If $\sigma(n)$ is odd, then np^l is half-Zumkeller if and only if $p \leq \sigma(n)$ and l is odd.*

Proof. (i). Let n be a practical number. Since $\sigma(n)$ is even, n is half-Zumkeller by Proposition 42. By Proposition 33, np^l is half-Zumkeller.

(ii). $\sigma(n)$ is odd. If np^l is half-Zumkeller, then np^l is Zumkeller by Fact 26. By Theorem 12, $p \leq \sigma(n)$ and l is odd. If $p \leq \sigma(n)$ and l is odd, then np^l is practical by Fact 8. Since l is odd, $\sigma(np^l) = \sigma(n) \sum_{i=0}^l p^i$ is even. By (i), np^l is half-Zumkeller. ■

A result similar to Proposition 15 can also be shown for half-Zumkeller numbers.

Proposition 44 *Let n be an even integer and p be a prime with $(n, p) = 1$. Then the following conditions are equivalent:*

- (i) *np is half-Zumkeller.*
- (ii) *The set of all positive factors of n can be partitioned into two disjoint parts D_1 and D_2 such that n is in D_1 , $\frac{n}{2}$ is in D_2 , and $p(\sum_{d \in D_2} d - \sum_{d \in D_1} d)$ can be written as a sum of some factors of n minus a sum of the rest of factors of n .*
- (iii) *The set of all positive factors of n can be partitioned into two disjoint parts D_1 and D_2 such that n is in D_1 , $\frac{n}{2}$ is in D_2 , and $\frac{(p+1)(\sum_{d \in D_2} d - \sum_{d \in D_1} d)}{2}$ can be written as a sum of some elements in D_2 possibly minus a sum of some elements in D_1 .*
- (iv) *The set of all positive factors of n can be partitioned into four disjoint parts A_1, A_2, A_3 and A_4 such that n is in either A_3 or A_4 , $\frac{n}{2}$ is in either A_1 or A_2 , and $(p+1) \sum_{d \in A_1} d + (p-1) \sum_{d \in A_2} d = (p+1) \sum_{d \in A_3} d + (p-1) \sum_{d \in A_4} d$.*

Proof. By Remark 27, np is half-Zumkeller if and only if the set D_0 of all positive factors of n can be partitioned into $D_0 = D_1 \cup D_2$ and $D_0 = D_3 \cup D_4$ such that

$$p\left(\sum_{d \in D_2} d - \sum_{d \in D_1} d\right) + \left(\sum_{d \in D_3} d - \sum_{d \in D_4} d\right) = 0,$$

and n is in D_2 and $\frac{n}{2}$ is in D_1 . The rest of the proof follows along the lines of the proof of Proposition 15. ■

Analogous to Fact 17 we have,

Fact 45 $2 \times 5^2 \times 7^2 \times p$ is half-Zumkeller for $p = 11, 13$.

Proof. In the proof that $2 \times 5^2 \times 7^2 \times p$ is Zumkeller for $p = 11, 13$. the partition of the set of all positive factors of $n = 2 \times 5^2 \times 7^2 = 2450$ into D_1 and D_2 satisfies the conditions that $n = 2450$ is in D_1 and $\frac{n}{2} = 1225$ is in D_2 in addition to condition (iii) in Proposition 15. So it satisfies condition (iii) in Proposition 44 and $2 \times 5^2 \times 7^2 \times p$ is half-Zumkeller for $p = 11, 13$. ■

We also have a result similar to Proposition 18 for half-Zumkeller numbers.

Proposition 46 If $1 = a_1 < a_2 < \dots < a_k = n$ are all factors of an even number n with $a_{i+1} \leq 2a_i$ for all i and $\sigma(n)$ is even, then n is half-Zumkeller.

Proof. Note that in the proof of Proposition 18, $a_k = n$ and $a_{k-1} = \frac{n}{2}$ have different signs. So we get a Zumkeller partition of n such that $a_k = n$ and $a_{k-1} = \frac{n}{2}$ are in distinct parts. By remark 27, n is half-Zumkeller. ■

As in remark 19, using Proposition 46, one can easily show that $n!$ is half-Zumkeller.

Our next result is similar to Remark 32. It gives more necessary conditions for a number to be an even Zumkeller number but not half-Zumkeller.

Proposition 47 Let n be an even Zumkeller with the prime factorization $2^k p_1^{k_1} p_2^{k_2} \dots p_m^{k_m}$, where $2 < p_1 < \dots < p_m$. If n is not half-Zumkeller, then there exists i , $1 \leq i \leq m$ such that $p_i > \sigma(2^k \dots p_{i-1}^{k_{i-1}}) + 1$. If j is the smallest such i , then $k_1, \dots, k_{j-1}$ must be even and $j \leq m - 1$.

Proof. Since n is Zumkeller, $\sigma(n)$ is even. If $p_i \leq \sigma(2^k \dots p_{i-1}^{k_{i-1}}) + 1$ holds for every i , $1 \leq i \leq m$, then n is practical. By Theorem 43 (i), n is half-Zumkeller. Therefore, there exists i , $1 \leq i \leq m$ such that $p_i > \sigma(2^k \dots p_{i-1}^{k_{i-1}}) + 1$. Let j be smallest such i . Then $2^k p_1^{k_1} \dots p_{j-1}^{k_{j-1}}$ is a practical number. If one of $k_1, \dots, k_{j-1}$ is odd, then $\sigma(2^k p_1^{k_1} \dots p_{j-1}^{k_{j-1}})$ is even. By Theorem 43, $2^k p_1^{k_1} \dots p_{j-1}^{k_{j-1}}$ is half-Zumkeller. By Proposition 33, $2^k p_1^{k_1} p_2^{k_2} \dots p_m^{k_m}$ is half-Zumkeller. This contradicts the hypothesis. So $k_1, \dots, k_{j-1}$ must be even. If $j = m$, then $2^k p_1^{k_1} \dots p_{m-1}^{k_{m-1}}$ is practical and $\sigma(2^k p_1^{k_1} \dots p_{m-1}^{k_{m-1}})$ is odd and

$p_m > \sigma(2^k p_1^{k_1} \cdots p_{m-1}^{k_{m-1}}) + 1$, then by Theorem 12(ii), $2^k p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$ is not Zumkeller and this contradicts the hypothesis that n is Zumkeller. $\blacksquare$

Next we show that Conjecture 2 is true for $n < 7.233498900 \times 10^9$. One can also show that it is true for $n = 7.233498900 \times 10^9$. But we omit the details.

Proposition 48 *Let n be even and Zumkeller. If n is not half-Zumkeller, then $n \geq 7.233498900 \times 10^9$.*

Proof. Let the prime factorization of n be $2^k p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$ with $2 < p_1 < \cdots < p_m$. By Proposition 47, there exists i , $1 \leq i \leq m$ such that $p_i > \sigma(2^k \cdots p_{i-1}^{k_{i-1}}) + 1$. Let j be the smallest such i . Then $k_1, \dots, k_{j-1}$ must be even and $j \leq m - 1$. We will use this j several times in the proof.

We will discuss the cases $k = 1, 2$ and $k \geq 3$ below.

Case 1. $k = 1$. In this case, $n = 2p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$.

Subcase 1(a): $m \geq 7$.

If $p_1 \neq 3$, then $p_1 \geq 5$.

If $7 \leq m \leq 10$, then $p_1 = 5$ and $p_2 = 7$. Since if $p_1 \geq 7$, then by Fact 1,

$$\frac{\sigma(n)}{n} < (1 + \frac{1}{2}) \prod_{i=1}^m \frac{p_i}{p_i - 1} < \frac{3}{2} \cdot \frac{7}{6} \cdot \frac{11}{10} \cdot \frac{13}{12} \cdot \frac{17}{16} \cdot \frac{19}{18} \cdot \frac{23}{22} \cdot \frac{29}{28} \cdot \frac{31}{30} \cdot \frac{37}{36} \cdot \frac{41}{40} < 3.$$

By Proposition 28, n is half-Zumkeller and this is a contradiction to our assumption. So $p_1 = 5$ when $m \leq 10$. If $p_2 > 7$, then $p_2 \geq 11$ and

$$\frac{\sigma(n)}{n} < (1 + \frac{1}{2}) \prod_{i=1}^m \frac{p_i}{p_i - 1} < \frac{3}{2} \cdot \frac{5}{4} \cdot \frac{11}{10} \cdot \frac{13}{12} \cdot \frac{17}{16} \cdot \frac{19}{18} \cdot \frac{23}{22} \cdot \frac{29}{28} \cdot \frac{31}{30} \cdot \frac{37}{36} \cdot \frac{41}{40} < 3.$$

By Proposition 28, n is half-Zumkeller and this is a contradiction to our assumption. So $p_2 = 7$ when $m \leq 10$ and $p_1 = 5$.

By Fact 23, $2 \cdot 5 \cdot 7$, $2 \cdot 5^2 \cdot 7$ and $2 \cdot 5 \cdot 7^2$ are half-Zumkeller numbers. By Proposition 36, $2 \cdot 5^{\text{odd}} \cdot 7^{\text{odd}}$, $2 \cdot 5^{\text{odd}} \cdot 7^2$, and $2 \cdot 5^2 \cdot 7^{\text{odd}}$ are half-Zumkeller numbers. By Fact 45, $2 \cdot 5^2 \cdot 7^2 \cdot p$ is half-Zumkeller for $p \leq 13$. Recall that $m \geq 7$, we have that

$$n \geq \min\{2 \cdot 5^2 \cdot 7^2 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31, 2 \cdot 5^4 \cdot 7^2 \cdot 11 \cdot 13 \cdot 17 \cdot 19 \cdot 23\} > 1.6 \times 10^{10}.$$

If $m \geq 11$, then $n \geq 2 \cdot 5 \cdot 7 \cdot 11 \cdot 13 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 37 \cdot 41 > 10^{14}$.

If $p_1 = 3$, then $2 \leq j \leq m-1$ (recall the meaning of j defined in the beginning of the proof) and recall that $k_1, \dots, k_{j-1}$ must be even. So

$$\begin{aligned}
n &> 2p_1^{k_1} \cdots p_{j-1}^{k_{j-1}} \sigma(2p_1^{k_1} \cdots p_{j-1}^{k_{j-1}})^{m-j+1} \\
&\geq 2p_1^2 \cdots p_{j-1}^2 \sigma(2p_1^2 \cdots p_{j-1}^2)^{m-j+1} \\
&\geq 2 \cdot 3^{2(j-1)} \cdot [(1+2)(1+3+3^2)^{j-1}]^{m-j+1} \\
&\geq 2 \cdot 3^{2(j-1)} (3 \cdot 13^{j-1})^{m-j+1} \\
&= 2 \cdot 3^{m+j-1} \cdot 13^{(j-1)(m-j+1)} \\
&= 2 \cdot 3^{m+j-1+(j-1)(m-j+1) \log_3 13}.
\end{aligned}$$

Since the exponent in the above expression is a quadratic function of j with negative coefficient of j , it reaches the minimum at either $j = 2$ or $j = m-1$ (recall that $2 \leq j \leq m-1$). By comparing the function values at these two places, we get that it has the minimum value when $j = 2$, so

$$n > 2 \cdot 3^{m+1} \cdot 13^{m-1} \geq 2 \cdot 3^8 \cdot 13^6 > 6.3 \times 10^{10}.$$

Subcase 1(b): $m \leq 6$.

If $p_1 \geq 5$, then by Fact 1,

$$\frac{\sigma(n)}{n} < (1 + \frac{1}{2}) \prod_{i=1}^m (1 + \frac{1}{p_i - 1}) < \frac{3}{2} \cdot \frac{5}{4} \cdot \frac{7}{6} \cdot \frac{11}{10} \cdot \frac{13}{12} \cdot \frac{17}{16} \cdot \frac{19}{18} < 3.$$

By Proposition 28, n is half-Zumkeller and this is a contradiction to our assumption. So

$$p_1 = 3.$$

So $j \geq 2$. Then by Proposition 47, $m \geq j+1 \geq 3$. By Proposition 30,

$$\frac{\sigma(n)}{n} \geq \frac{10}{3}.$$

If $p_2 > \sigma(2 \cdot p_1^{k_1}) + 1 \geq \sigma(2 \cdot 3^2) + 1 = 40$, then $p_2 \geq 41$ and by Fact 1

$$\frac{\sigma(n)}{n} < (1 + \frac{1}{2}) \prod_{i=1}^m \frac{p_i}{p_i - 1} < \frac{3}{2} \cdot (\frac{41}{40})^5 < \frac{10}{3}.$$

So

$$p_2 \leq \sigma(2 \cdot p_1^{k_1}) + 1.$$

So $j \geq 3$ and $m \geq j+1 \geq 4$.

If $p_3 > \sigma(2 \cdot 3^{k_1} 5^{k_2}) + 1 \geq \sigma(2 \cdot 3^2 5^2) + 1 = 1210$, then

$$\frac{\sigma(n)}{n} < \frac{3}{2} \cdot \frac{3}{2} \cdot \frac{5}{4} \cdot \left(\frac{1211}{1210}\right)^4 < \frac{10}{3}.$$

So $p_3 \leq \sigma(2 \cdot 3^2 5^2) + 1$. Therefore $j \geq 4$ and $m \geq j + 1 \geq 5$ by Proposition 47. If $p_4 > \sigma(2 \cdot 3^{k_1} 5^{k_2} p_3^{k_3}) + 1 \geq \sigma(2 \cdot 3^2 5^2 7^2) + 1 = 68914$, then

$$\frac{\sigma(n)}{n} < \frac{3}{2} \cdot \frac{3}{2} \cdot \frac{5}{4} \cdot \frac{7}{6} \cdot \left(\frac{68915}{68914}\right)^3 < \frac{10}{3}.$$

So $p_4 \leq \sigma(2 \cdot 3^{k_1} 5^{k_2} p_3^{k_3}) + 1$. Therefore, $j \geq 5$ and by Proposition 47, $m \geq j + 1 \geq 6$. Since $m = 6$, then $j = 5$ and

$$p_5 > \sigma(2 \cdot 3^{k_1} 5^{k_2} p_3^{k_3} p_4^{k_4}) + 1 \geq \sigma(2 \cdot 3^2 5^2 7^2 11^2) + 1 = 9165430.$$

So

$$n > 2 \cdot 3^2 5^2 7^2 11^2 (9165431)^2 > 2 \times 10^{22}.$$

Case 2. $k = 2$. In this case, $n = 2^2 p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$.

Subcase 2(a): $m \geq 7$.

If $p_1 \leq 7 \leq \sigma(2^2) + 1$, then $2 \leq j \leq m - 1$. Recall that all $k_1, \dots, k_{j-1}$ are even. So

$$\begin{aligned} n &> 2^2 p_1^{k_1} \cdots p_{j-1}^{k_{j-1}} \sigma(2^2 p_1^{k_1} \cdots p_{j-1}^{k_{j-1}})^{m-j+1} \\ &\geq 2^2 p_1^2 \cdots p_{j-1}^2 \sigma(2^2 p_1^2 \cdots p_{j-1}^2)^{m-j+1} \\ &\geq 2^2 \cdot 3^{2(j-1)} (7 \cdot 13^{j-1})^{m-j+1} \\ &= 2^2 \cdot 3^{2j-2} \cdot 7^{m-j+1} \cdot 13^{(j-1)(m-j+1)} \\ &= 2^2 \cdot 3^{2j-2+(m-j+1)\log_3 7 + (j-1)(m-j+1)\log_3 13}. \end{aligned}$$

Since the exponent in the above expression is a quadratic function of i with negative coefficient of i , it reaches the minimum at either $j = 2$ or $j = m - 1$. By comparing the function values at these two places, we get that it has the minimum value when $j = 2$, so

$$n > 2^2 \cdot 3^2 \cdot 7^{m-1} \cdot 13^{m-1} \geq 36 \cdot 7^6 \cdot 13^6 > 2 \times 10^{13}.$$

If $p_1 \geq 11$, then $m \geq 14$. Since if $m \leq 13$ then

$$\frac{\sigma(n)}{n} < \left(1 + \frac{1}{2} + \frac{1}{4}\right) \prod_{i=1}^m \frac{p_i}{p_i - 1} < \frac{7}{4} \cdot \frac{11}{10} \cdot \frac{13}{12} \cdot \frac{17}{16} \cdot \frac{19}{18} \cdot \frac{23}{22} \cdot \frac{29}{28} \cdot \frac{31}{30} \cdot \frac{37}{36} \cdot \frac{41}{40} \cdot \frac{43}{42} \cdot \frac{47}{46} \cdot \frac{53}{52} \cdot \frac{59}{58} < 3.$$

By Proposition 28, n half-Zumkeller and this is a contradiction to our assumption. So $m \geq 14$ if $p_1 \geq 11$. In this case,

$$n \geq 2^2 \cdot 11 \cdot 13 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 37 \cdot 41 \cdot 43 \cdot 47 \cdot 53 \cdot 59 \cdot 61 > 2.2 \times 10^{21}.$$

Subcase 2(b): $m \leq 6$.

If $m \leq 3$, then $p_1 = 3$ since otherwise, $\frac{\sigma(n)}{n} < (1 + \frac{1}{2} + \frac{1}{2^2}) \cdot \frac{5}{4} \cdot \frac{7}{6} \cdot \frac{11}{10} = \frac{7}{4} \cdot \frac{5}{4} \cdot \frac{7}{6} \cdot \frac{11}{10} < 3$ and n is half-Zumkeller by Proposition 28, this contradicts our assumption. So $j \geq 2$ and $m \geq j + 1 \geq 3$ by Proposition 47. Since $p_1 = 3$, by Proposition 30, $\frac{\sigma(n)}{n} \geq \frac{10}{3}$.

If $m = 3$: If $p_2 > \sigma(2^2 p_1^{k_1}) + 1 \geq \sigma(2^2 3^2) + 1 = 92$, then $\frac{\sigma(n)}{n} < \frac{7}{4} \cdot \frac{3}{2} \cdot \frac{97}{96} \cdot \frac{101}{100} < \frac{10}{3}$. Therefore, $p_2 \leq \sigma(2^2 p_1^{k_1}) + 1$. So $j \geq 3$ and $m \geq j + 1 \geq 4$.

What left is to consider $m = 4$ or 5 or 6 . If $p_1 > \sigma(2^2) + 1 = 8$, then $p_1 \geq 11$ and

$$\frac{\sigma(n)}{n} < \frac{7}{4} \cdot \frac{11}{10} \cdot \frac{13}{12} \cdot \frac{17}{16} \cdot \frac{19}{18} \cdot \frac{23}{22} \cdot \frac{29}{28} < 3.$$

So $p_1 \leq \sigma(2^2) + 1$. Therefore, $2 \leq j \leq m - 1$.

If $j = 2$, then $p_2 > \sigma(2^2 p_1^{k_1}) + 1 \geq \sigma(2^2 3^2) + 1 = 92$, so $p_2 \geq 97$ and

$$\frac{\sigma(n)}{n} < \frac{7}{4} \cdot \frac{3}{2} \cdot \frac{97}{96} \cdot \frac{101}{100} \cdot \frac{103}{102} \cdot \frac{107}{106} < 3.$$

Therefore, $j \geq 3$.

If $j = 3$, then $p_3 > \sigma(2^2 p_1^{k_1} p_2^{k_2}) + 1 \geq \sigma(2^2 3^2 5^2) + 1 = 2822$ and $m \geq j + 1 \geq 4$. So $p_3 \geq 2833$, $p_4 \geq 2837$, and

$$n \geq 2^2 3^2 5^2 \cdot 2833 \cdot 2837 = 7.202859300 \times 10^9.$$

It can be shown that $2^2 3^2 5^2 \cdot 2833 \cdot 2837$ is half-Zumkeller. We omit the details.

If $j = 4$, then $p_4 > \sigma(2^2 p_1^{k_1} p_2^{k_2} p_3^{k_3}) + 1 \geq \sigma(2^2 3^2 5^2 7^2) + 1 = 160798$ and $m \geq j + 1 \geq 5$. So

$$n > 2^2 3^2 5^2 7^2 160799^2 > 1.1 \times 10^{15}.$$

If $j = 5$ (only for $m = 6$), then $p_5 > \sigma(2^2 p_1^{k_1} p_2^{k_2} p_3^{k_3} p_4^{k_4}) + 1 \geq \sigma(2^2 3^2 5^2 7^2 11^2) + 1 = 21386002$. So

$$n > 2^2 3^2 5^2 7^2 11^2 21386003^2 > 2.4 \times 10^{21}.$$

Case 3. $k \geq 3$. In this case, $n = 2^3 p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$.

Subcase 3a: $m \geq 7$.

If $p_1 > \sigma(2^k) + 1 \geq \sigma(2^3) + 1 = 16$, then $p_1 \geq 17$ and

$$n \geq 2^3 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 37 \cdot 41 > 8.1 \times 10^{10}.$$

If $p_1 \leq \sigma(2^k) + 1$, then $2 \leq j \leq m - 1$. Recall that $k_1, \dots, k_{j-1}$ must be even. So

$$\begin{aligned} n &> 2^k p_1^{k_1} \cdots p_{j-1}^{k_{j-1}} (\sigma(2^k p_1^{k_1} \cdots p_{j-1}^{k_{j-1}}))^{m-j+1} \\ &\geq 2^3 3^{2j-2} (15 \cdot 13^{j-1})^{m-j+1} \\ &\geq 2^3 3^2 15^{m-1} 13^{m-1} \geq 2^3 3^2 15^6 13^6 > 3.9 \times 10^{15}. \end{aligned}$$

Subcase 3b: $m \leq 6$.

If $p_1 > 7$, then $p_1 \geq 11$ and

$$\frac{\sigma(n)}{n} < 2 \cdot \frac{11}{10} \cdot \frac{13}{12} \cdot \frac{17}{16} \cdot \frac{19}{18} \cdot \frac{23}{22} \cdot \frac{29}{28} < 3.$$

So $p_1 \leq 7$. So $j \geq 2$ and $m \geq j + 1 \geq 3$ by Proposition 47.

We will discuss the cases $p_1 = 3$ or $5 \leq p_1 \leq 7$ separately as below.

Subcase 3b1: $p_1 = 3$.

If $p_2 > \sigma(2^3 3^{k_1}) + 1 \geq \sigma(2^3 3^2) + 1 = 196$, then $p_2 \geq 197$ and

$$\frac{\sigma(n)}{n} < 2 \cdot \frac{3}{2} \cdot \left(\frac{197}{196}\right)^5 < \frac{10}{3}.$$

By Proposition 30, n is half-Zumkeller and this contradicts our assumption. So $p_2 \leq \sigma(2^3 3^{k_1}) + 1$. So $j \geq 3$ and $m \geq j + 1 \geq 4$ by Proposition 47.

So we can assume that $4 \leq m \leq 6$. Recall that $3 \leq j \leq m - 1$ and $k_1, \dots, k_{j-1}$ must be even. So

$$\begin{aligned} n &> 2^k p_1^{k_1} \cdots p_{j-1}^{k_{j-1}} (\sigma(2^k p_1^{k_1} \cdots p_{j-1}^{k_{j-1}}))^{m-j+1} \\ &\geq 2^3 3^2 (5^2)^{j-2} (15 \cdot 13 \cdot 31^{j-2})^{m-j+1} \\ &\geq 2^3 3^2 5^{2(m-3)} 15^2 13^2 31^{2(m-3)} (j = m - 1) \\ &\geq 2^3 3^2 5^2 15^2 13^2 31^2 > 6.5 \times 10^{10}. \end{aligned}$$

Subcase 3b2: $5 \leq p_1 \leq 7$.

If $p_2 > \sigma(2^3 p_1^{k_1}) + 1 \geq \sigma(2^3 5^2) + 1 = 466$, then $p_2 \geq 467$ and

$$\frac{\sigma(n)}{n} < 2 \cdot \frac{5}{4} \cdot \left(\frac{467}{466}\right)^5 < 3.$$

Since n is not half-Zumkeller, then this is impossible by Proposition 28. So $p_2 \leq \sigma(2^3 p_1^{k_1}) + 1$. So $j \geq 3$ and $m \geq j + 1 \geq 4$ by Proposition 47.

So we can assume that $4 \leq m \leq 6$. Recall that $3 \leq j \leq m - 1$ and $k_1, \dots, k_{i-1}$ must be even. So

$$n > 2^k p_1^{k_1} \cdots p_{j-1}^{k_{j-1}} (\sigma(2^k p_1^{k_1} \cdots p_{j-1}^{k_{j-1}}))^{m-j+1}$$

$$\begin{aligned}
&\geq 2^3 5^{2j-2} (15 \cdot 31^{j-1})^{m-j+1} \\
&\geq 2^3 5^4 15^{m-2} 31^{2(m-2)} \quad (j = 3) \\
&\geq 2^3 5^4 15^2 31^4 > 1.03 \times 10^{12}. \blacksquare
\end{aligned}$$

5 Problems

The following problems need further study.

- 1.[1] Is every even Zumkeller number half-Zumkeller?
2. What are the odd Zumkeller numbers?
3. What are the odd half-Zumkeller numbers?
4. Does the set of Zumkeller numbers have density? Note that this set is a subset of abundant numbers and the set of abundant numbers has density between 0.2474 and 0.2480 [2].

References

- [1] S. Clark, J. Dalzell, J. Holliday, D. Leach, M. Liatti and M. Walsh, Zumkeller numbers, presented in the Mathematical Abundance Conference at Illinois State University on April 18th, 2008.
- [2] Marc Degeglise, Bounds for the density of abundant integers, *Experimental Mathematics* 7 (1998): 137-143.
- [3] A. Galletti and K.P.S. Bhaskara Rao, A new proof of the Egyptian fraction theorem and great numbers, manuscript.
- [4] A. K. Srinivasan, Practical numbers, *Current Science* 17 (1948): 179-180, MR0027799.
- [5] B. M. Stewart, Sums of distinct divisors, *American Journal of Mathematics* 76 (1954): 779-785.